

## Psychosis, Schizophrenia and Schizoaffective Disorder

Duffy, F.H. et al. Neurophysiological differences between patients clinically at high risk for schizophrenia and neurotypical controls—first steps in development of a biomarker. BMC Med. 2015 Nov 2;13,276

Light, G.A., Swerdlow, N.R. Future clinical uses of neurophysiological biomarkers to predict and monitor treatment response in schizophrenia. Ann N Y Acad Sci. 2015 May; 1344,105-119

Moeini, M, Khaleghi, A, Amiri, N, Niknam, Z. Quantitative electroencephalogram (QEEG) Spectral Analysis of Patients with Schizoaffective Disorder compared to Normal Subjects. Iran j Psychiatry. 2014 Oct;9(4),216-221

Fuggetta, G et al. Quantitative electroencephalography as a biomarker for proneness toward developing psychosis. Schizophr Res. 2014 Mar;153(1-3),68-77.

Kim, J.W. et al. Diagnostic utility of quantitative EEG in unmedicated schizophrenia. Neurosci Lett. 2015 Mar 4;589,126-131

Beth Israel Deaconess Medical Center. Researchers ID, Treat Faulty Circuitry Underlying Symptoms of Schizophrenia. Science Daily January 30, 2019

Kwon, D. The Cerebellum is Your Little Brain – and it Does Some Pretty Big Things. Scientific America January 18, 2019